

Perspectives On The POINT-DIPOLE Approximation:
for the
Prospective INTUITIVE-CHEMISTS' Approaches

S.Aravamudhan, Department of Chemistry, North Eastern Hill University, NEHU Campus SHILLONG 793022
Meghalaya INDIA

**ADDRESSES FOR SOME OF THIS AUTHOR'S WEBPAGES WHERE MATERIALS RELEVANT TO THE
TOPIC HERE ARE DOCUMENTED.**

WebSite URL (I) : <http://saravamudhan.tripod.com>

Explaining the trends of Nuclear Shielding caused by Magnetic moments related to Susceptibilities

Home The Foreground and the current trends of research activities of Dr.S.Aravamudhan

Page 1 The 2nd Alpine Conference on Solid State NMR, Chamonix Mont-Blanc, France, 9-13th Sept.2001

Page 2 Contribution to ISMAR-CA '98 TU Berlin, Aug.1998

Page 3 The Summing procedure and the volume elements with the Benzene Molecular perspectives

Page 4 Shielding- and Demagnetization-factor Calculations

Page 5 A Graphical Depiction of the Results on the Calculation of Demagnetization Factors

Page 6 Explanation and illustration of Induced Field Calculations at Points Outside the Specimen Close to it

Page 7 Any Spill Overs from OR to the nanostructures in materials?

Page 8 Pictorial Depiction of Semimicro Volume Elements Graphical Plots of Calculations

Page 9 Contribution to the XIV International Biophysics Congress, Argentina April 27-May 1, 2002

Page 10 About the Author/Curriculum Vitae/List of Contributions and Publications

WebSite URL (II): <http://geocities.com/amudhan20012000/Confview.html>

WebSite URL (III): http://www.geocities.com/amudhan20012000/Conf_View_Poster_Sheets_Certificates.html

WebSite URL (IV) : http://geocities.com/amudhan_nehu/graphpresent.html

ABSTRACT

A dipole by definition [**SHEET 2**] consists of two unlike poles but equal in magnitude (pole strength) separated by a finite distance '**d**'. When the point (where this dipole's effect is) being considered is located at such large distances '**R**' from the dipole that '**d**' is negligible compared to '**R**' [$d \ll R$ (as in **SHEET 8 and 9**)], the point-dipole approximation is said to be valid. When '**R**' becomes nearly equal (comparable in magnitude) to '**d**', then the point dipole-approximation would not be reliable.


Then, by [1] hypothetically considering [**SHEET 10**] each of the poles to be made up of 'n' number of equally divided pole strengths and [2] placing each of this divided poles separated from the correspondingly divided unlike-pole, and, further, [3] this distance of separation being equal to '**d**' = d/n , a set of 'n' smaller dipoles can be constructed all these dipoles being centered at the same Electrical Centre of Gravity as the original undivided, single dipole. The value of 'n' can be so chosen that the distances of separation '**d**' is negligible compared to the value **R**. This would enable the point dipole approximation to be valid for each of the divided, smaller dipoles and their effect at a point at distance '**R**' can be calculated. Since all the 'n' number of divided dipoles are identical their contributions will all be the same and having calculated the effect of one of the dipoles, the effect of 'n' dipoles can be obtained by a multiplication by 'n' [**A critical question 'n' or 'n²**]. This should yield the effect of the single dipole within the frame work of point dipole approximation. Does this mean that dividing and multiplying by 'n' [**or 'n²**] [**effectively the multiplication factor is '1' or 'n'**] makes it a valid description that which was originally not reliable? This procedure would be providing an argument to say the same numerical values are reliable even though at the outset this result could have been discarded as unreliable.

Since this is an argument which can lead a chemist astray while trying to interpret the results, it is intended in this poster presentation to address this question in greater detail so as to enable a confident handling of the point-dipole approximation which is inevitably the beginners' approach in chemical contexts.


INDEX to the Display SHEETS

SHEET 2 and SHEET 3 : Definition of dipole, Dipole Moment **SHEET 4 and SHEET 5** : Charge cloud descriptions, Circulation and movement of Charges and associated CURRENT FLOW. Flowing Current as source of Magnetic Dipole Moment **SHEET 5 and SHEET 6**: Rotational Motions of Electrons (Orbital and Spin) and spin of Nuclei: The associated angular momenta and the Magnetic Moments.\.: as is well evidenced in the description of the Magnetic Resonance Phenomena. **SHEET 7**: Such Magnetic Moments, Occurrence, Origin and consequences in Physical Chemistry **SHEET 8**: A Consideration of the POINT-DIPOLE Approximation: Criteria for its validity **SHEET 9,10**: How to make the Point-Dipole approximation more valid while the determining factors become critically uncompromisable. **SHEET 10,11** : General indications of Induced Fields Due to circulation of electrons causing demagnetizing and Shielding effects **SHEET 12**: Summary and Conclusions

These set of equally spaced parallel set of lines indicate the presence of homogeneous




Direction of movement of +ve pole in external field



Direction of movement of -ve pole in external magnetic field

The two poles, equal in magnitude and opposite in sign, are not held fixed in a non-flexible and hard framework. Hence in external fields the two poles can be moving independent of each other and hence the inter-pole distance d is not a constant. Thus the two poles cannot be said to be forming a “dipole” with a well defined value for the “dipole moment”.



In the above set of illustration the pair of poles are held by non-flexible and hard FRAME-work and the inter-pole distance d cannot alter in presence of the external field. When a movement is required the two poles move together with fixed distance between them. When no translational motion can provide the required change to the minimum potential energy situation, the minimization occurs by a rotation of the line (imaginary) connecting the two poles. This is said to align the “dipole” or the “Dipole moment” gets disposed along the direction of the external field.

Image (1) of the 2 images for this display







Image (2)

Fig. 3.10

Fig. 3.11



Strength of north pole = m
 Strength of south pole = m

MAGNETIC DIPOLE MOMENT = $\vec{\mu}$

$\vec{\mu}$ POINTS FROM S to N INSIDE
 THE BAR MAGNET


magnitude $|\vec{\mu}| = 2lm$

All the magnetic field effect representations can be done by simply considering the magnetic dipole moment vector $\vec{\mu}$ instead of picturing the bar magnet with its north pole and south pole.

Thus $\vec{\mu}$ represents the dipole characteristics.

While explaining the presence of a “**dipole moment**” by considering the poles which, depending on the context, the poles may be the “electrical (positive and negative) charges” or the [“magnetic charges”] magnetic “north” and “south” poles. In either case a dipole moment can be defined: in the former case it would be electrical dipole moment and in the latter case it would be magnetic dipole moment

In the case of flowing electrical charges, the current thus flowing can cause a magnetic field, which can be associated with the flow characteristics and the amount of the electrical charges. The magnetic field thus arising can be calculated and the direction of the field determined. This can be the attribute “dipole moment” for the flowing charge systems.



Flowing positive charges

In each of the above cases a magnetic moment can be attributed for the charge flow. In the case of SHAPE 3 the charge cloud is depicted. The total charge cloud can be considered as small, charged, volume elements inside the charge cloud. These volume elements each can be attributed a small dipole moment and the total dipole moment associated with currents in the charge cloud can be obtained by a vector addition of the individual elemental dipole moments.

IF THE PARTICLE UNDERGOING A ROTATIONAL MOTION ALSO HAS AN ELECTRICAL CHARGE, THEN THE ROTATIONAL MOTION OF THE CHARGE — i.e., a flowing charge — WILL PRODUCE A MAGNETIC FIELD.

IT IS WELL KNOWN THAT IF A CURRENT 'I' FLOWS THROUGH A CIRCULAR COIL AND IF THE AREA OF THE CIRCLE IS 'A', THEN THE MAGNETIC FIELD PRODUCED IS GIVEN BY $\mu = IA$ and the magnetic field is at the center of the circle.

